

Structurally specific thermal fluctuations identify functional sites for DNA transcription

G. Kalosakas, K. Ø. Rasmussen, and A. R. Bishop

*Theoretical Division and Center for Nonlinear Studies,
Los Alamos National Laboratory, Los Alamos, New Mexico 87545*

C. H. Choi and A. Usheva

*Endocrinology, Beth Israel Deaconess Medical Center, Department of Medicine,
Harvard Medical School, 99 Brookline Ave., Boston, Massachusetts 02215*

We report results showing that thermally-induced openings of double stranded DNA coincide with the location of functionally relevant sites for transcription. Investigating both viral and bacterial DNA gene promoter segments, we found that the most probable opening occurs at the transcription start site. Minor openings appear to be related to other regulatory sites. Our results suggest that coherent thermal fluctuations play an important role in the initiation of transcription. Essential elements of the dynamics, in addition to sequence specificity, are nonlinearity and entropy, provided by local base-pair constraints.

PACS numbers: 87.15.Aa, 87.15.Ya, 87.15.He

One of the most challenging subjects in biophysics is the relation between biomolecular motions and function [1]. We present an example suggesting that functionality arises from structurally coherent dynamics, with essential ingredients of: *sequence-specificity*, *nonlinearity* and *entropy*; the nonlinearity from *local constraints* is crucial. In particular, remarkably successful comparisons of numerical simulations of a minimal model (see below) of transverse dynamics for gene promoter DNA segments with *in vitro* transcriptional experiments, shows that the combination of *all* the above mentioned components controls coherent “bubble” fluctuational openings of base-pairs around specific sites of promoter DNA. Remarkably, the prominent opening occurs at the transcription initiation site, while minor openings coincide with regulatory sites at which transcription factors and other assisting proteins are bound. These results demonstrate the importance of the sequence structure, not simply as a static and passive element, but to *provide the template for specific coherent fluctuations determining function*. These coherent structures constitute a *colored* spatio-temporal stochastic environment. This is an example of the importance of a (dynamic) landscape of substates [1].

We have used a microscopic model proposed by Peyrard and Bishop [2] to describe the dynamics of the openings of double stranded DNA. This model focuses only on the most relevant degrees of freedom, namely the transverse stretching of the hydrogen bonds connecting complementary bases in the opposite strands of the double helix. Its reduced character, involving a small number of variables, makes it suitable for simulations over relatively long times and appropriate for gathering sufficient statistics. Subsequent key improvements [3] succeeded in reproducing the abrupt (first order) character of the observed DNA denaturation transition. The potential energy of this model reads [3]

$$V = \sum_n \left[D_n (e^{-a_n y_n} - 1)^2 + \frac{k}{2} (1 + \rho \exp[-\beta(y_n + y_{n-1})]) (y_n - y_{n-1})^2 \right] \quad (1)$$

Here the sum is over all the base-pairs of the DNA and y_n denotes the displacement from the equilibrium position of the relative distance between the bases within the n^{th} base-pair, divided over $\sqrt{2}$. The Morse potential (other similar potentials can also be used) in the first term provides the effective interactions between complementary bases; it represents both the attraction due to the hydrogen bonds forming the base-pairs and the repulsion of the negatively charged phosphates in the backbone of the two strands screened by the surrounding solvent. Beyond the exact details of this interaction, an important issue is a correct description of the nonlinearities. The parameters D_n and a_n of the on-site potential distinguish between the two possible combinations of bases, i.e. adenine-thymine (A-T) or guanine-cytosine (G-C), at site n , depending on the particular sequence. The second term in the total potential energy represents the stacking interaction potential between adjacent base-pairs. Here the nonlinear inter-site coupling, given by the exponential term that effectively modifies a harmonic spring constant, is essential for representing *local constraints* in nucleotide motions, which result in long-range cooperative effects [3]. As in elastic materials [4, 5], it controls lattice vibrations, yielding accurate entropic terms [6]: the stiffening of the coupling in the compact state compared to that in the open state leads to an abrupt entropy-driven transition [3]. Physically, the constraint describes the change of the next-neighbor stacking interaction due to the distortion of the hydrogen bonds connecting a base-pair, mediated through the redistribution of the electrons on

FIG. 2: Auto-radiography of $[^{32}P]$ -labeled reverse transcripts after separation by gel electrophoresis (lanes 3). Arrows indicate the direction of specific transcription started from the initiation site +1 in all cases. Lanes 1 indicate base position markers obtained by chemical sequencing. (a) P5 promoter, (b) AdMLP promoter, and (c) T7 promoter. Lane 4 in (a) shows elimination of the transcription for the mutated P5 promoter, where the nucleotides at +2 and +3 have been changed from AT to GC.

displacements y_n of the base-pairs at each site n and the following $m - 1$ (m varying from 1 to 19) base-pairs. If the openings at *all* these m subsequent sites are greater than a threshold value y_{th} (varying from one tenth to few A) we assign a contribution to the opening event at the n^{th} base-pair of the sequence. The obtained opening probabilities along the studied DNA segments for bubble sizes of $m = 10$ base-pairs and thresholds $y_{th} = 1.0$ and $1.5 A$ for accepting an opening are presented in Fig. 3. (We recall that the real openings are equal to $y\sqrt{2}$). In all these cases the most probable openings are located at the transcription start site +1 (see Fig. 2). Furthermore, in the viral cases the other distinct openings seem to be related to known regulatory sites: in P5 the opening at the A/T rich region between -40 and -35 corresponds to the binding site of the transcription factor Yin Yang 1 [15], while in AdMLP the second higher opening is close to the binding site of the TATA-box binding protein [16], that is necessary for transcription. As can be seen from Fig. 3, openings of such large widths and amplitudes are rare events in our microscopic simulations, therefore requiring sufficient statistics.

We stress that similar local sequences do *not* exhibit the same opening probabilities; equal size segments of relatively weakly bound A/T pairs in different parts of the promoter show very different statistics (compare for example the region around -30 with that around +1 in P5). Furthermore, the larger openings do not occur in regions with longer A/T stretches, as might be intuitively




FIG. 3: Logarithm of the probability P for the occurrence of an opening of 10 base-pair width and amplitude of more than $2.1 A$ (thick solid line) or $1.4 A$ (dotted line) starting at a particular site n of the DNA fragment, as a function of n , for the (a) P5 promoter, (b) AdMLP promoter, and (c) T7 promoter.

anticipated because of the weaker bonding. Effective long range cooperativity (from the nonlinear inter-site potential in equation (1)) and competing localization lengths due to the spatial disorder and the nonlinearity are responsible for this high specificity: in general, length scale competition in nonlinear systems is known [17] to lead to complex spatio-temporal (dynamic landscape) behavior. The sensitivity of the cooperative/competing phe-




FIG. 4: Logarithm of the probability P for the occurrence of an opening of 10 base-pair width and amplitude of more than 2.1 Å (thick solid line) or 1.4 Å (dotted line) starting at a particular site n of the DNA fragment, as a function of n for the mutated P5 promoter (see text).




FIG. 5: Logarithm of the probability P for the occurrence of an opening of 10 base-pair width and amplitude of more than 2.1 Å (thick solid line) or 1.4 Å (dotted line) starting at a particular site n of the DNA fragment, as a function of n for the P5 promoter, by linearizing the stacking interaction term of the potential (i.e. setting $\rho = 0$ in potential energy (1)).

nomena, affected even by a *single* base-pair modification, enhances the predictive power of our model; a small mutation of the sequence (at a specific location) is sufficient to completely eliminate bubble formation at the transcription initiation site. In Fig. 4 we show numerical calculations of the opening probabilities for a mutated P5 promoter, where nucleotides +2 and +3 have been changed from AT to GC. This mutation completely eliminates the opening at the previous transcriptional start site, in agreement with no transcriptional events occurring in the corresponding experiment (see Fig. 2a, lane 4).

We emphasize that, as in previous applications of this model, the nonlinear inter-site coupling is crucial for its success [3, 10]. For example, as can be seen in Fig. 5, linearizing the stacking interaction term ($\rho = 0$) results in

very modified statistics for the openings of the P5 promoter, changes the position of the peaks along the sequence, and eliminates the successful comparison with the experimental observations. The nonlinear inter-site coupling constitutes a minimal representation of the local stacking constraint between neighboring base-pairs. As in more general situations of displacive structural phase transitions[4, 5], such local constraints can lead to long-range “elastic” interactions and macroscopic cooperativity.

In summary, our model and simulations suggest that structurally specific coherent thermal fluctuations identify locations in the DNA sequences where the RNA polymerase initiates transcription. Further, we find indications that the thermal fluctuations also help in recruiting other protein complexes participating in the transcriptional process, by separating the DNA double strand at specific locations. These bubbles precede protein binding and their possible role is limited to the very initial steps of the transcription. This suggests that DNA, through structurally specific dynamics, participates in directing its own transcription.

We would like to thank H. Frauenfelder, P. Fenimore, and J.A. Krumhansl for valuable discussions. This research was supported by the US DoE, under contract W-7405-ENG-36 and NIH.

-
- [1] H. Frauenfelder, S.G. Sligar, and P.G. Wolynes, *Science* **254**, 1598 (1991).
 - [2] M. Peyrard and A.R. Bishop, *Phys. Rev. Lett.* **62**, 2755 (1989).
 - [3] T. Dauxois, M. Peyrard, and A.R. Bishop, *Phys. Rev. E* **47**, R44 (1993).
 - [4] J.R. Morris and R.J. Gooding, *Phys. Rev. Lett.* **65**, 1769 (1990); *Phys. Rev. B* **43**, 6057 (1991).
 - [5] W.C. Kerr, A.M. Hawthorne, R.J. Gooding, A.R. Bishop, and J.A. Krumhansl, *Phys. Rev. B* **45**, 7036 (1992).
 - [6] The importance of entropic terms for enhancing cooperativity and providing distinct transitions, similarly to the case of the model we used, has also been stressed recently in a different context, viz. β -hairpin folding models. See C. Guo, H. Levine, and D.A. Kessler, *Phys. Rev. Lett.* **84**, 3490 (2000).
 - [7] T. Dauxois and M. Peyrard, *Phys. Rev. E* **51**, 4027 (1995).
 - [8] We note that these large-amplitude bubbles (breathers) are not traveling, in contrast to mobile low-amplitude nonlinear excitations used to account for hydrogen exchange experiments in, e.g., S.W. Englander, N.R. Kallenbach, A.J. Heeger, J.A. Krumhansl, and S. Litwin, *Proc. Natl. Acad. Sci. USA* **77**, 7222 (1980).
 - [9] A. Campa and A. Giansanti, *Phys. Rev. E* **58**, 3585 (1998).
 - [10] D. Cule and T. Hwa, *Phys. Rev. Lett.* **79**, 2375 (1997).
 - [11] S. Komineas, G. Kalosakas, and A.R. Bishop, *Phys. Rev. E* **65**, 061905 (2002).
 - [12] M. Peyrard and J. Farago, *Physica A* **288**, 199 (2000).

- [13] G. Kalosakas, K.Ø. Rasmussen, and A.R. Bishop, J. Chem. Phys. **118**, 3731 (2003); to be published in Synthetic Metals.
- [14] C.H. Choi et al., submitted to Nucleic Acids Research.
- [15] A. Usheva and T. Shenk, Cell **76**, 1115 (1994).
- [16] M.F. Concino, R.F. Lee, J.P. Merryweather, and R. Weinmann, Nucleic Acids Research **12**, 7423 (1984).
- [17] A. Sanchez and A.R. Bishop, SIAM Review **40**, 579 (1998).

This figure "Fig2.gif" is available in "gif" format from:

<http://arXiv.org/ps/cond-mat/0309157v2>